

MANAGEMENT of COUGH

Jed Gorden MD

University of Washington

Vietnam Lecture Series 2001

Three Categories of Cough

- Acute Cough = < 3 Weeks Duration
- Subacute Cough = 3 – 8 Weeks Duration
- Chronic Cough = > 8 Weeks Duration

Acute Cough < 3 Weeks

Differential Diagnosis

Acute Cough

- Upper Respiratory Tract infections:
 - Viral syndromes, sinusitis viral / bacterial
- Allergies
- Exacerbation of Chronic Obstructive Pulmonary Disease (COPD)
- Left Ventricular Heart Failure
- Pneumonia
- Foreign Body Aspiration

Common Cold/Viral Rhinosinusitis

- Presentation:
- **Symptoms** – Nasal Passages
 - Rhinorrhea, Sneezing, Nasal obstruction, Post nasal drip
- **Signs** - +/- Fever, +/- throat irritation, normal chest auscultation
- **Diagnostic** – No Laboratory or X-ray

Common Cold/Viral Rhinosinusitis

- **Treatment**

- Antihistamine (H1) + Pseudoephedrine

OR

- Naproxen

Treatment Failure

Viral Rhinosinusitis

VS

Bacterial Rhinosinusitis

Viral vs. Bacterial Rhinosinusitis

- **Viral**

- Most Common
- Treat empirically

- **Bacterial**

- Less Common
- Treat in cases of treatment failure
- Treat for set criteria

Criteria Bacterial Rhinosinusitis

- Treatment failure

+

Two of the following signs or symptoms

1. Maxillary Tooth Ache
2. Purulent Nasal Discharge
3. Abnormal Sinus Trans-illumination
4. Discolored Nasal Discharge

Treatment

- Antihistamine + Pseudoephedrine

+

Oxymetazoline (Afrin)

+

Antibiotics against Haemophilis influenza
and Streptococcus pneumonia

(Bactrim TMP/Sulfa or Amoxicillin)

Subacute Cough 3-8 Weeks

Subacute Cough Differential Diagnosis

- Postinfectious
- Bacterial Sinusitis
- Asthma

Post Infectious Cough

- A cough that begins with an acute respiratory tract infection and is not complicated* by pneumonia

*Not complicated = Normal lung exam
normal chest X-ray

Post Infectious Cough

- Post Infectious cough will resolve without treatment
- Cause = Postnasal drip or Tracheobronchitis

Indications For Chest X-ray

- Abnormal auscultatory lung exam

Chest X-ray: Management

Treat Abnormality

- Infiltrate = Pneumonia = Antibiotics
- Cardiomegaly/Pulmonary Edema = Heart Failure
- Normal Chest X-ray Consider Empiric Therapy for Asthma

Chronic Cough > 8 Weeks

Chronic Cough

Differential Diagnosis

- Post Nasal Drip (Nose and Sinus Conditions)
 - Gastroesophageal Reflux Disease
 - Chronic Bronchitis from Tobacco
- Chronic Obstructive Pulmonary Disease
 - Left Ventricular Heart Failure
 - Lung Cancer
 - Tuberculosis
 - Asthma

Patients Who Present With
Chronic Cough Should Receive a
Chest X-ray When Possible

Chest X-ray and Differential Diagnosis

- **Normal X-ray**
 - **Post Nasal drip**
 - **Reflux Disease**
 - **Asthma**
 - **Chronic Bronchitis**
- **Abnormal X-ray**
 - **Tuberculosis**
 - **COPD**
 - **Heart Failure**
 - **Lung Cancer**

Specific Causes of Cough Focus

- Asthma
- Chronic Obstructive Pulmonary Disease
(COPD)

Asthma

- Asthma is a Chronic Inflammatory Disorder of the Airway
- Activation of the Immune System = Airway Hyperresponsiveness + Airflow Limitation
- Airflow Limitation is Reversible

Asthma

- Asthma is Present in all Age Groups
- Asthma Affects Men and Women Equally

Asthma

Signs and Symptoms

- Signs and Symptoms Vary from Patient to Patient as well as being Dynamic over time
- Classic Symptoms: Wheezing
 - Shortness of Breath
 - Cough
 - Chest Tightness

Asthma Precipitants

- Many Nonspecific Precipitants Provoke Asthma Symptoms and the Need for Medication
 - Respiratory Infections
 - Exercise
 - GI Reflux
 - Stress
 - Weather Changes

Asthma Treatment

- Based on Symptom Severity
 - “Step” Approach

“Step Up” Meds = Poor Symptom Control

“Step Down” Meds = Good Control

Classification Asthma Severity

Mild Intermittent	Symptoms < 2X Week Exacerbations Brief Rare	Night Symptom < 2 X Month
Mild Persistent	Symptoms > 2X Week < 1X Day Exacerbations Rare +/- Limit Activity	Night Symptom > 2 X Month
Moderate Persistent	Symptoms Daily Symptoms Limit Activity Exacerbations > 2 per Week	Night Symptom > 1 X Week
Severe Persistent	Continuous Symptoms Limited Exercise Tolerance Frequent Exacerbations	Night Symptom Frequent

Asthma: Stepwise Management

	Long Term Control	Quick Relief
Step 4 Severe Persistent	Inhaled Steroid (High Dose) + Long Acting Bronchodilator (B2 Agonist or Theophylline) + Oral Steroids	Short Acting Bronchodilator Inhaled B2 Agonist
Step 3 Moderate Persistent	Inhaled Steroid (Med Dose) Or Long Acting Bronchodilator (B2 Agonist or Theophylline) + Inhaled Steroids	Short Acting Bronchodilator Inhaled B2 Agonist

Asthma: Stepwise Management

	Long Term Control	Rapid Relief
Step 2 Mild Persistent	Low Dose Inhaled Steroids	Short Acting Bronchodilator B2 Agonist
Step 1 Mild Intermittent	No Daily Medications	Short Acting Bronchodilator B2 Agonist

Asthma: Medications

The Dose of Medication that Reaches
the Lung is Dependant On:

- Delivery Device
 - Drug Dose
- Patient Technique

Inhaled B2 Agonist

- Most Effective Drug for the Treatment of Acute Bronchospasm and Prevention of Exercise Induced Asthma
- Example: Albuterol, Proventil, Ventolin

B2 Agonist

- Target: Selective for Bronchodilation
- Toxicity: Tachycardia, Palpitations, Tremor
Extreme overuse May = Hypokalemia

Inhaled Corticosteroids

- Target: Suppress Inflammation, Minimize Airway Hyperresponsiveness
- Toxicity: Rare Stunt Growth in Children, Dermal Thinning

Theophylline (Methylxanthine)

- Target: Smooth Muscle Dilation of the Bronchial Tree, Anti-Inflammatory, Mucociliary Clearance
- Toxicity: Nausea, Nervousness, Headache, Insomnia, Vomiting, Tachycardia, Tremor, Seizures

Oral Steroids

- Target: Most Effective Therapy for Decreasing Inflammation and Airway Hyperresponsiveness
- Toxicity: Glucose Intolerance, Weight Gain, Hypertension, Osteoporosis

Asthma Diagnosis

Asthma Diagnosis

- History
- Physical Exam
- Clinical Suspicion/Response Empiric Trial
- Pulmonary Function Testing

Pulmonary Function Testing

Lung Volume

Airflow Rates

Ability to Transfer Gas Across the Alveolar
Capillary Membrane

Flow Rates Define Asthma

- Forced Vital Capacity (FVC) = Volume of Gas that can be expelled from the lungs After Maximal Inspiration
- Forced Expiratory Volume in 1 Sec (FEV₁) = Volume of Gas Expelled in the First Second of the FVC Maneuver

Flow Rate Compromise

- FEV₁/FVC 75% Mild Obstruction
- FEV₁/FVC 50-75% Moderate Obstruction
- FEV₁/FVC <50% Severe Obstruction
 - REVERSIBILITY
- Increase 12% and 200 cc in FEV₁ OR
- Increase 15% and 200 cc in FVC

Provocative Testing: Methacholine Challenge

- Positive Test = Decrease in FEV1 of at Least 20% at a dose of 16mg/ml or less
- A negative Test has a Negative Predictive Value For Asthma of 95%

Chronic Obstructive Pulmonary Disease (COPD)

COPD: Definition

- Airflow Obstruction From Chronic Bronchitis or Emphysema; Airflow obstruction is Progressive, may be accompanied by Airway Hyperreactivity and May be Partially Reversible

COPD: Terms

- Chronic Bronchitis= Cough for Three Months in any 2 Successive Years without other Cause
- Emphysema = Pathologic Diagnosis Describing Airspace Destruction

COPD: Risk Factors

- **SMOKING/TOBACCO**
- Genetic Alpha1 Antitrypsin Deficiency
(Less Than 3% of Cases)
- Environmental/Occupational Exposure

Natural History of COPD

- CHART HERE

COPD: Diagnosis

- History
- Physical Examination
- Laboratory and Spirometry

COPD: Patterns of Advanced Disease

Pink Puffer (Emphysema)

- Dyspnea
- Age > 50
- Rare Cough
- Thin/Weight Loss
- Quiet Auscultatory Exam
- No Peripheral edema

Blue Bloater (Bronchitis)

- Chronic Cough/Productive
 - Age > 40
- Dyspnea Mild
- Over Weight
- Cyanotic
- Chest + Rhonchi/Wheezes

COPD: Patterns of Advanced Disease

Pink Puffer (Emphysema)

- Normal Hematocrit
- PaO₂ Reduced
- PaCO₂ Normal/Reduced
- X-Ray = Hyperinflation

Blue Bloater (Bronchitis)

- Hematocrit Elevated
- PaCO₂ Elevated
- X-Ray Increased Markings
(Dirty X-Ray)

COPD: History

- Smoking 20 Cigarettes/Day > 20 Years
- >40 Years Old
- Dyspnea > 50 years Old
- Cough

COPD: Physical Exam

- Prolonged Expiration
- Expiratory Wheezing

Severe COPD

- Over distention of Lungs/ Increased A-P Diameter
- Decreased Heart Sounds
- Decreased Breath Sounds
- Pursed Lip Breathing
- Use of Accessory Muscles in Breathing

COPD: Chest X-ray

- Lung Distention = Long Narrow Heart Shadow
Flat Diaphragm
- Bullae = Radiolucent Areas > 1 cm in Diameter
(Caution Bullae can be Confused with Pneumothorax)

COPD: Pulmonary Function Test

- Stage I = $FEV_1 > 50\%$ Predicted
- Stage II = $FEV_1 35-49\%$ Predicted
- Stage III = $FEV_1 < 35\%$ Predicted

- $FEV_1 < .75$ L 1 Year Mortality = 30%
10 Year Mortality = 95%

COPD: Treatment

- **Stop Smoking**
- Smoking Cessation is Challenging:
 - Without Intervention 5% Success
 - With Intensive Intervention 22% Success at 5 Years
(US Lung Health Study)

COPD: Therapy Goals

- Induce Bronchodilation
- Decrease Inflammatory Response

COPD: Medication

- Bronchodilators = Beta2-Agonists = Albuterol
- Anticholinergic Agents = Ipratropium (Atrovent)
- Theophylline
- Anti-Inflammatory Therapy = Corticosteroides

COPD: Management

- Mild Variable Symptoms

B₂-Agonist 1-2 Puffs Every 2-6 Hours

- Mild – Moderate Continued Symptoms

Ipratropium 2-6 Puffs Every 6 Hours

+

B₂-Agonist

COPD: Management

- Inadequate Response to Ipratropium + B₂ Agonist
- Add Sustained Release Theophylline 200-400mg
2x/day
- Nocturnal Symptoms Theophylline 400-800mg

COPD: Management

- If Continued Poor Control

Corticosteroids – Prednisone 40mg 1x/day
For 10-14 Days*

If No Improvement Stop Abruptly


If Improvement Considered Inhaled Steroides

COPD: Management

- COPD Therapy Holds Many Similarities to Asthma
Important to Note Response to Treatment in COPD
Considerably Less than Response in Asthma
- Patient Population older less Tolerant and More
Sensitive to Drug Side Effects

Cough Summary

- Diverse Differential Diagnosis Involving Multiple Organ Systems
- Therapeutic Approach Requires Knowledge of Epidemiology and Symptom Complex
- Patient Care Requires Therapeutic Trial Which May Require Re-evaluation if Inadequate



CASE I

- 33 Year Old Male Presents For Care Complaining of 10 Days of Nonproductive Cough. Patient states Cough Syndrome was Preceded by Sinus Congestion, Muscle Aches and Fatigue.
- Patient has No Past Medical History
- Patient Lives With His Wife and 6 Year old Boy. The Child has been Irritable with Rhinorea

CASE I

- Physical Exam

HR 87 BP 140/70 RR 14 T 37.6

Sclera are Injected Bilaterally

Left Nares limited Air Flow

Sinuses Nontender with Good Transillumination

Lungs Clear to Auscultation

Cardiac Regular Without Murmur

CASE 1

- What is The Differential Diagnosis
- What Studies do You Need For Diagnosis
- What is Your Treatment Plan

CASE 2

- A 65 year Old Male Presents For Care, He is Complaining of Cough Worse in The Morning, Shortness of Breath and Increasing Dyspnea on Exertion. The Cough is Minimally Productive of Sputum
- No Past Medical History
- Patient Lives With His Wife He Smokes 1 Pack of Cigarettes/day for 40 Years

CASE 2

- Physical Exam
- BP 165/88 HR 75 RR 18 T 37.6
- Sinuses Non Tender No Rhinorea
- Lungs Crackles at Bases
- Chest Increased AP Diameter
- Cardiac Distant Heart Sounds

Case 2

- What is The Differential Diagnosis
- What Studies are Necessary For Diagnosis
- What is The Treatment Plan

CASE 3

- A 23 Year Old Female Present for Care Complaining of > 1 month of Cough The Cough is Not Productive of Sputum. The Cough is Worse When She Exercises or is Exposed to Cold Air. The Cough is Associated with Shortness of Breath
- The Patient Has No Past medical problems
- The Patient Is a University Student she Lives alone in a Dormitory She Does Not Smoke

CASE 3

- Physical Exam

HR 70 BP 140/60 RR 12 T 37.6

Sinuses Non Tender No Rhinorea

Lungs Diffuse Musical Wheezes With a
Prolonged Expiratory Phase

Cardiac Regular Without Murmur

Case 3

- What is The Differential Diagnosis
- What Studies Do You Need for diagnosis
- What is Your Treatment Plan